

**For Veterinary Use Only**

# Cloptic®

**Eye Ointment**

**Veterinary Antibacterial**

## **COMPOSITION**

It contains Cloxacillin Benzathine, equivalent to 835 mg Cloxacillin, in 5.0 g syringe. Cloptic Eye Ointment is a white, sterilized, and greasy ointment.

## **PHARMACOLOGICAL PROPERTIES**

Cloptic is a smooth, sterilized cream intended to ophthalmic comprising cloxacillin use in a retard formulation. Cloxacillin 6 is a semisynthetic penicillin derived from Amino-penicillanic acid. Staphylococci sp. and Bacillus sp. are main bacteria amongst sensitive bacteria, and because it is resistant to penicillinases, it is also effective against staphylococci strains resistant to penicillin G. Cloptic is a locally effective antibiotic, which is not combined into bloodstream when administered into eyes. After administering Cloptic its density in tear is over the effective MIC levels for 2-3 days, thus it provides an efficient therapy and it allows to a simple administration. Following administration it is excreted from the body through tear.

## **AREA OF USE/INDICATIONS**

Cloptic is used in therapies for eye infections caused by sensitive bacteria including Staphylococcus sp. (including beta-lactamase producing strains), Bacillus sp. in cattle, sheep, horse, dog and cats.

## **USAGE AND DOSAGE**

Cloptic is administered to eyes locally for only treatments of eye infections. After opening administration tube pomade is administered as a stripe to conjunctival space by opening lower eyelid.

### **Practice dose scale**

<b><u>Animal Species</u></b>	<b><u>Dose</u></b>
Cattle - Horse	A stripe around 5 to 10 cm per eye
Sheep	A stripe around 5 cm per eye
Cat - Dog	A stripe around 2 cm per eye

In cases with one sick eye to both of those eyes should be applied medicine against cross contamination risk. In order to prevent infection risk first to that sick eye should be applied medicine.

## **UNDESIRE/SIDE EFFECTS**

Penicillin may cause allergic reactions (type 1 allergy) in sensitive humans and animals.

## **DRUG INTERACTIONS**

Avoid using together with tetracycline.

## **LEGAL PURIFICATION TIME**

***Withholding period (WHP): For Meat and Milk: is "0" zero day.***

**CONTRAINDICATIONS**

Avoid administrating in animals with known hypersensitivity to penicillin and cephalosporin.

**GENERAL WARNINGS**

Consult your veterinary surgeon before using and in case an undesired effect is observed. Keep out of reach of children. Avoid using products with damaged packages.

**SPECIAL WARNINGS FOR APPLIER AND VETERINARY SURGEON**

The contact of the drug in the syringe with human skin may cause allergic reactions in people who are sensitive to penicillin. Use surgical gloves during administration. It is not recommended for the appliers who are sensitive to penicillinic products to use **Cloptic** suspension. In case of symptoms such as skin rash following contact through any routes, consult a physician immediately. In case of swelling of face, lips and eyes or respiratory distress, seek for medical assistance immediately.

**STORAGE CONDITIONS AND SHELF LIFE:**

Store under 25 °C, and protect from light. Shelf life is 2 years (24 months).

**COMMERCIAL PRESENTATION FORM :**

It is presented as 5 g poliethylen syringes in cardboard boxes for a single package, and as 4, 12, and 40 pieces in one more cardboard boxes.

**PLACE AND CONDITIONS OF SALE**

Sold with Veterinary Surgeon's prescription in pharmacies and veterinary surgeries.

**APPROVAL DATE OF PACKAGE INSERT:** 07.01.2009

**MARKETING AUTHORIZATION DATE AND NO.** 29.12.2008 / 20 – 097

**NAME AND ADDRESS OF MARKETING AUTHORIZATION OWNER:**

**Alke Sağlık Ürünleri San. ve Tic. A.Ş**

Dolayoba, Çınardere Mh. 3. Petek Sk.

No: 18, 34896 Pendik-İSTANBUL

**NAME AND ADDRESS OF MANUFACTURER COMPANY:**

**Alke Sağlık Ürünleri San. ve Tic. A.Ş**

Merkez Organize Sanayi Bölgesi 2. Kısım 10. Cadde No:7 Tokat